

Dynamical transitions in Markovian exciton transport

Hoda Hossein-Nejad¹, Alexandra Olaya-Castro², Francesca Fassioli^{2,3}, and Gregory D Scholes³

¹Department of Physics, University of Toronto, 60 St. George Street, Toronto, Ontario M5S 1A7, Canada,

²Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK

³Lash-Miller Chemical Laboratories, Institute for Optical Sciences and Centre for Quantum Information and Quantum Control, University of Toronto, 80 St. George Street, Toronto, Ontario, M5S 3H6 Canada

E-mail: h.hossein-nejad@ucl.ac.uk

PACS numbers: 05.70.Ln, 03.65.Yz, 42.50.Lc

Abstract.

We investigate the counting statistics of non-thermal equilibrium, or rare, quantum jump trajectories associated to Markovian exciton transport. By applying the large deviation theory to open quantum system dynamics, we are able to uncover dynamical transitions, or crossovers, in the statistics of such rare trajectories. We find that the crossovers are activated by electronic coherences, and occur in systems with a minimum of three interacting molecules. The formalism thus reveals subtle consequences of non-equilibrium thermal fluctuations on the steady-state statistics of systems exhibiting Markovian transport.

1. Introduction

Full counting statistics (FCS) is a powerful mathematical tool that enables the characterization of a counting process by systematic computation of the generator of its statistics. The technique was originally developed for theoretical modelling of mesoscopic charge transport [1, 2], where current fluctuations can provide vital information on the nature of the transport mechanism [3, 4, 5, 6, 7, 8]. FCS of a classical, or an open quantum system, at thermal equilibrium can be obtained by identifying the generating function of the statistics from the corresponding master equations [9, 10, 11]. However, due to the environmental perturbations, non-equilibrium fluctuations can easily arise in open quantum systems. This raises the question that how are the counting statistics altered by such perturbations, and whether a systematic theory of non-equilibrium FCS for open quantum systems can be formulated.

The difficulty in addressing this problem lies in the different nature of the two influences: thermal fluctuations of the environment are of a classical character and are described via the tools of non-equilibrium thermodynamics. The interaction of the quantum system with the environment is described via the theory of open quantum systems [12]. There has been a historical distinction between the two approaches, rendering the study of their mutual influence a non-trivial one.

An important connection between the two descriptions was provided by Garrahan and Lesanovsky in a recent contribution [13]. The link between the two descriptions was provided through the well-known mathematical theory of large deviation principle [14]. In statistical mechanics, large deviation principle provides a framework to deduce thermodynamic quantities from sub-ensembles of particles in the configuration space. In the theory of open quantum systems, however, it serves as a tool for statistical analysis of rare (i.e.: non-equilibrium) quantum-jump trajectories associated to Markovian dynamics. A quantum-jump trajectory is a realization of the system's dynamics subjected to particular bath-induced measurements at stochastic intervals, and represents a sequence of configurations in the time domain. By incorporating a coupling field, or a *dynamical order parameter*, that quantifies perturbations from equilibrium, the large deviation approach grants access to the complete statistics of the trajectories, both in and out of equilibrium. At equilibrium, large deviation theory recovers the results of the full counting statistics as expected. Garrahan and Lesanovsky coined the term the *s-ensemble* theory to a technique of mapping out the out of equilibrium trajectories of a quantum master equation via the large deviation approach. This terminology will also be used here.

In the context of open quantum systems, the *s-ensemble* approach allows the identification of statistical transitions in the time domain, thereby providing novel physical insight on the dynamics of quantum systems subjected to Markovian noise. Dynamical transitions have so far been predicted in photon counting [13, 15, 16, 17], electron counting [18], and quadrature measurements [19], suggesting the possibility of their existence in a host of other systems. The significance of transitions in the statistics of quantum processes, or the means by which they may be observed, are not always clear and tend to be problem-specific. It is evident, however, that such novel dynamical features cannot be identified from the analysis of the density matrix operator alone.

In this article we apply the *s-ensemble* theory to investigate the statistics of rare trajectories of quantum evolutions describing excitation energy transport in a molecular complex. Assuming strong electronic interactions, and weak system-environment coupling, exciton relaxation can be described by Lindblad type evolution [12]. The exciton relaxation process results in incoherent transfer between electronic eigenstates, reaching a steady-state where the average population of the eigenstates obeys the thermal Boltzmann distribution. At a microscopic level, however, transfer of excitation still occurs in

a dynamical manner in the steady-state. For instance for a system of two eigenstates at thermal equilibrium, detailed balance dictates that $k_{1 \rightarrow 2}P_1 = k_{2 \rightarrow 1}P_2$ where $k_{i \rightarrow j}$ is the rate of energy transfer from the state i to the state j and P_i is the thermally-equilibrated population of the state i . Out of equilibrium fluctuations result in deviations from the detailed balance. Therefore, in the present context, the s -ensemble theory serves as a mathematical tool to investigate changes in the counting statistics as the steady-state deviates from thermal equilibrium. We demonstrate that despite the classical nature of the transitions, the statistics of the trajectories are not, in general, Poissonian. We observe temperature-dependent dynamical transitions for systems containing a minimum of three sites, and properties such as bunching and anti-bunching that are often associated with non-classical correlations.

This article is organized as follows: in section 2 we introduce the conceptual and mathematical framework of the large deviation principle and discuss its application to counting statistics. In section 3 the Markovian Lindblad master equation and the s -ensemble theory are introduced. In section 4 we apply the s -ensemble theory to investigate the statistics of rare trajectories in systems of two, three, four and seven sites and present numerical results that demonstrate the dynamical transitions in the statistics of trajectories. The electronic and vibrational parameters of the Fenna-Matthews-Olson (FMO) photosynthetic complex are used in the computation of the numerical results, as the Lindblad master equation considered in this article has been found to make a reasonable prediction of the steady-state of the complex. Section 5 summarizes the main conclusions of the article.

2. Large deviation principle and counting statistics

A counting process can be decomposed into a number of trajectories where in each trajectory the process occurs at different stochastic intervals [20]. In the context of quantum jumps, the counting process is the number of jumps observed after a given time t . Let us denote the probability of observing K jumps after a time t by $P_t(K)$, where $\sum_{K=0}^{\infty} P_t(K) = 1$, and $\{0 \leq P_t(K) \leq 1\}$. For a counting process that satisfies the large deviation theory, $P_t(K)$ acquires a large deviation form for large t . That is

$$P_t(K) \simeq e^{-t\varphi(k)} \quad (1)$$

where the function $\varphi(k)$ is known as the rate function [14], and $k = K/t$ is the jump rate. For a simple (unimodal) counting process, $\varphi(k)$ has a local minima corresponding to the most probable jump rate after a long observation time. The generating function of a given probability distribution is define to be

$$Z_t(s) = \sum_{K=0}^{\infty} P_t(K) e^{-sK} \quad (2)$$

where s is a conjugate variable to K and is known as the *activation parameter*. If the counting process has a large deviation form, the generating function can be written as

$$Z_t(s) \simeq e^{t\theta(s)}. \quad (3)$$

$Z_t(s)$ is now referred to as the large deviation function, while $\theta(s)$ is the cumulant generating function. The cumulant generating function and the rate function are related via a Legendre transform,

$$\theta(s) = -\min_k [\varphi(k) + ks] \quad (4)$$

The average and the standard deviation of the number of jumps at equilibrium can be deduced from the cumulant generating function, and are given by

$$\langle k \rangle = -\theta'(0) \quad (5)$$

$$\langle k^2 \rangle - \langle k \rangle^2 = \theta''(0). \quad (6)$$

Similarly, higher derivatives of $\theta(s)$ evaluated at $s = 0$, are the higher order statistical cumulants. Full counting statistics (or the equilibrium statistics) is therefore recovered by looking at the derivatives of the cumulant generating function at $s = 0$. The applicability of the large deviation approach to non-equilibrium processes becomes evident when the behaviour of the cumulant generating function away from $s = 0$ is considered. To illustrate this let us construct an associated stochastic process defined by the probabilities [21]

$$q_t(K, s) = \frac{1}{Z_t(s)} P_t(K) e^{-sK} \quad (7)$$

For $s = 0$ we recover the original counting process. For $s < 0$ ($s > 0$) the exponential factor amplifies (lowers) the probabilities of rare events. The coupling field s , thus provides a systematic means by which rare events can be accessed: unlikely events of the counting process $\{P_t(K)\}$, correspond to typical events of the associated process $\{q_t(K, s)\}$. Rare events can thus be accessed without generating a large number of typical events and relying on the statistics of the original counting process. In non-equilibrium thermodynamics, the coupling field s is interpreted as a means by which the system is perturbed out of equilibrium. *The influence of non-equilibrium perturbations are therefore deviations from the thermodynamic probabilities.* A rare event corresponds to a particular trajectory of the system that has been significantly perturbed out of equilibrium.

In statistical mechanics the counting process is the number of microstate configurations accessible to the system under certain constraints, making $Z_t(s)$ the partition function. Analogously, for an open quantum system, $Z_t(s)$ is a measure of the number of trajectories that can participate in the dynamics of a particular non-equilibrium configuration. One can also identify $\varphi(k)$ as the entropy, and $\theta(s)$ as the free energy. Extrema of $\varphi(k)$ therefore correspond to the equilibrium configurations, or the typical trajectories. The variable s is (minus) the chemical potential $s \equiv -\mu$ [21]. Free energy serves as an order parameter: singularities in free energy predict first-order

thermodynamic phase transitions, or dynamical transitions in the space of trajectories of a quantum system.

3. *s*-ensemble theory and electronic energy transfer

In this section we apply the large deviation theory to investigate the trajectories of a particular open quantum system, modelled by the Markovian Lindblad master equation. The system of interest consists of a number of molecules (sites) coupled via electronic couplings and weakly interacting with their environment. Each molecule is modelled as a two-level system in its electronic coordinate, where the two levels correspond to the ground and the first excited electronic states. The electronic degrees of freedom are assumed to be diagonally coupled to a bosonic bath of harmonic oscillators. We assume that there is one excitation in the system that can be coherently shared between multiple molecules (a molecular exciton). The Hamiltonian of this system is given by

$$H = - \sum_{ij} J_{ij} |i\rangle\langle j| + \sum_{i=1}^2 |i\rangle\langle i| \left[\epsilon + \frac{1}{2} \sum_k h_{ik} (b_k^\dagger + b_k) \right] + \sum_k \omega_k b_k^\dagger b_k \quad (8)$$

where J_{ij} is the electronic coupling between the ground state of site i and excited state of site j , the ket $|i\rangle$ represents the excited state of site i , h_{ik} is the electron-phonon coupling between site i and mode k of the bath, $\{b_k^\dagger, b_k\}$ are the raising and lowering operators for mode k of the bath, and ω_k are the bath frequencies.

Due to the assumption of weak system-bath interaction, the exciton states are delocalized and the influence of the environment is to induce jumps between them at stochastic intervals. In the steady-state the average population of the exciton states is given by the Boltzmann factor: $P_i = e^{-\beta\omega_i}/Z$, where P_i is the population of the exciton state with energy ω_i and Z is the partition function. From a microscopic standpoint, dynamic transfer of excitation still persists in the steady-state. For instance for a system of two sites, steady-state implies that $k_{1\rightarrow 2}P_1 = k_{2\rightarrow 1}P_2$, where $k_{i\rightarrow j}$ is the transfer rate from the eigenstate i to the eigenstate j . Moreover, the ratio of the transfer rates satisfies detailed balance: $k_{1\rightarrow 2} = k_{2\rightarrow 1}e^{-\beta\omega_{12}}$, where ω_{12} is the energy difference between the two states. $k_{1\rightarrow 2}P_1$ is also the average number of jumps per second between the two states: upward and downward jumps occur with equal frequency in the steady-state. If the environment is perturbed out of equilibrium, detailed balance no longer holds and the frequency of jumps deviates from the thermally equilibrated rate.

To compute the statistics of jumps in the steady-state, one can compute the cumulant generating function of this process directly from the Lindblad equation. The Lindblad equation as applied to EET, for weak system-bath coupling with secular approximation is given by [12]

$$\dot{\sigma}(t) = -i[H_s, \sigma(t)] + \sum_{\omega, m} \gamma(\omega) \left[A_m(\omega) \sigma(t) A_m^\dagger(\omega) - \frac{1}{2} \{A_m^\dagger A_m(\omega), \sigma(t)\} \right] \quad (9)$$

where the commutator describes the unitary evolution and the second term describes relaxation and dephasing in the eigenstates $\{|\alpha\rangle\}$ of the unperturbed system

Hamiltonian H_s (neglecting the Lamb-shift Hamiltonian). The Lindblad operators are given by $A_m(\omega) = \sum_{\omega=\epsilon_\alpha-\epsilon_{\alpha'}} c_m^*(\alpha)c_m(\alpha')|\alpha\rangle\langle\alpha'|$ where $\{\omega\}$ is the energy difference associated with single excitation eigenstates, and $c_m^*(\alpha)c_m(\alpha')$ is the *exciton interference* at site m . The factor $\gamma(\omega)$ determines the rate of energy transfer between two excitons and is related to the spectral density of the bath $J(\omega)$, via the expression $\gamma(\omega) = 2\pi J(|\omega|)|n(\omega)|$, where $n(\omega) = (e^{\beta\omega} - 1)^{-1}$ is the thermal occupation number. In the present treatment it is assumed that each molecule interacts with an identical and independent phonon bath. The dissipative term in the Lindblad equation describes the incoherent jumps between the exciton states as induced by the phonon bath. In the quantum jump approach the Hamiltonian evolution of the quantum system is interrupted by jumps at stochastic intervals. A sequence of stochastic jumps constitutes a trajectory of the system, and an averaging over a large number of trajectories recovers the reduced density matrix $\sigma(t)$ [22].

If K jumps are observed after a time t , the (unnormalized) reduced density matrix of the system prior to detection is given by the projection of the total density matrix onto the subspace of K events [9, 10]. We denote this projected density matrix by $\sigma^{(K)}(t)$. Conversely, the total density matrix can be written as an expansion of all projected density matrices, that is $\sigma(t) = \sum_{K=0}^{\infty} \sigma^{(K)}(t)$. The probability of observing K events after a time t is given by $P_t(K) = \text{Tr}[\sigma^{(K)}(t)]$. At this point for the sake of clarity we restrict the analysis to the statistics of jumps between two exciton states separated in energy by ω_1 . Note that the total system may still contain an arbitrary number of excitons. It can be shown that the Laplace transform of the projected density operator, defined by the expression $\sigma_s(t) = \sum_0^{\infty} \sigma^{(K)} e^{-sK}$ obeys the equation $\dot{\sigma}_s(t) = W_s[\sigma_s]$, where the superoperator W_s is defined to be

$$\begin{aligned} W_s[\sigma_s(t)] = & -i[H_s, \sigma_s(t)] + \sum_m \gamma(\omega_1) e^{-s} A_m(\omega_1) \sigma_s(t) A_m^\dagger(\omega_1) \\ & + \sum_m \left[\sum_{\omega \neq \omega_1} \gamma(\omega) A_m(\omega) \sigma_s(t) A_m^\dagger(\omega) - \frac{1}{2} \sum_{\omega} \gamma(\omega) \{A_m^\dagger(\omega) A_m(\omega), \sigma_s(t)\} \right]. \end{aligned} \quad (10)$$

Note that $\sigma_s(t)$ has the same structure as the variable $q_t(K, s)$ in Eq. (7). Typical (equilibrium) trajectories of $\sigma_s(t)$ thus correspond to rare trajectories of the original master equation. The ensemble of trajectories generated by Eq. (11) is known as the s -ensemble [13]. Physical time evolution is retrieved when the activation parameter is zero ($s = 0$), whereas $s \neq 0$ contains information with regards to rare trajectories of the system. $s < 0$ ($s > 0$) generates the active (inactive) trajectories where the number of jumps are significantly higher (lower) than average.

The Markovian nature of the Lindblad equation enables us to compute the statistical moments of the number of jumps in the steady-state via the large deviation principle. Large deviation theory states that for a Markov process, the largest real eigenvalue of the superoperator W_s is the cumulant generating function of the statistics [14]. Following the notation of the previous section, we denote this generator by $\theta(s)$. In the next section we compute the cumulant generating function and the associated statistical moments for a system of two, three, four and seven sites.

4. Results and discussion

Adopting the electronic and vibrational parameters of the FMO complex, we first consider two and three sites (chromophores) and study the statistics of trajectories that contain jumps between two given exciton states. We subsequently include further sites in the system and study the statistics of selected jumps. The coupling of the electronic system to the vibrational environment is described via the spectral density

$$J(\omega) = \frac{2E_r}{\pi} \frac{\omega\omega_c}{\omega^2 + \omega_c^2} \quad (11)$$

where $E_r = 35 \text{ cm}^{-1}$ is the reorganization energy $E_r = \int_0^\infty \frac{J(\omega)}{\omega} d\omega$, and $\omega_c = 150 \text{ cm}^{-1}$ is the cutoff frequency of the bath [23, 24]. We compute the matrix W_s and solve for the cumulant generating function $\theta(s)$. Figure 1 (a) and 1(b) illustrate the exciton states where two and three sites are included in the dynamics respectively. The two site model includes sites 1 and 2, and the three site model includes sites 1, 2 and 3. The site indices are the standard indices assigned to the chromophores in the FMO complex [25]. The exciton states are labelled via a greek letter and a number, where the greek letter represents the position of the exciton state in the energy ladder, and the number indicates the site that carries the maximum exciton amplitude. For instance $|\alpha : 1\rangle$ represents the lowest eigenstate, and indicates that this eigenstate is predominately localized on site 1. Due to significant exciton localization in the FMO protein, each exciton can be unambiguously assigned to a unique site index.

Figure 1 (c) and (d) illustrate the cumulant generating function versus the activation parameter s for different temperatures. The plots suggest that the two cases have similar statistics in the active phase, but begin to diverge close to $s = 0$. Figure 1 (e) illustrates the Mandel parameter for the case of two sites. Mandel parameter is an indicator of the nature of first-order statistical correlations in the system and is defined to be

$$Q(s) = \frac{\langle k^2 \rangle - \langle k \rangle^2}{\langle k \rangle} - 1 = -\frac{\theta''(s)}{\theta'(s)} - 1. \quad (12)$$

$Q = 0$ implies Poissonian statistics where probabilities of subsequent events are uncorrelated. $Q > 0$ ($Q < 0$) indicates super-Poissonian (sub-Poissonian) statistics where subsequent jumps exhibit first-order correlations.

For the case of two sites, the Mandel parameter is negative for all activations. This indicates sub-Poissonian correlations (anti-bunching) where a waiting period for re-excitation is required before the system can relax back to the lower state. As the activation parameter is increased, the Mandel parameter approaches zero (Poissonian limit). In this limit there are so few jumps that correlations vanish asymptotically.

It is striking that anti-bunching is observed even though the transitions are thermally activated, and the steady-state dynamics obey a classical rate equation. The classical nature of the correlations can be confirmed by considering the classical rate

equation characterized by the following dynamical matrix

$$W_{class}(s) = \begin{bmatrix} -\kappa & \Gamma e^{-s} \\ \kappa & -\Gamma \end{bmatrix}$$

where κ and Γ are the relevant equilibrium transfer rates, and $\Gamma = \kappa e^{-\beta\omega}$ as required by detailed balance. By diagonalizing $W_{class}(s)$, we arrive at the following expression for $Q(s)$

$$Q(s) = \frac{-2\kappa\Gamma}{(\kappa + \Gamma)^2 - 4\kappa\Gamma(1 - e^{-s})} e^{-s} < 0 \quad \forall s \quad (13)$$

Indeed, the same expression for $Q(s)$ is obtained if the full Lindblad equation with dephasing terms is instead considered. *The formalism thus captures the fluctuations of a quantum system due to its coupling to a non-equilibrium bath. For the system under consideration the nature of these fluctuations is entirely classical.*

The corresponding plot of the Mandel parameter for a system of three sites is shown in Fig. 1 (f). The trajectories exhibit a dynamical crossover in their statistics, i.e.: the correlations change from sub-Poissonian ($Q(s) < 0$) to super-Poissonian ($Q(s) > 0$) as the activation parameter is varied. The crossover point is temperature dependent and is closer to $s = 0$ at higher temperatures. Transitions close to $s = 0$ are more significant as they indicate that small perturbations from equilibrium can significantly change the nature of statistical correlation. Typical trajectories ($s = 0$) are bunched, and the Mandel parameter has a peak close to $s = 0$.

One can continue to add more sites to the model and investigate the statistics of a particular jump, or the collective statistics of a number of jumps. For instance the collective statistics of all jumps to the lowest eigenstate can be investigated. In general, beyond a few sites there are many possible jumps and there is no obvious advantageous collective counting strategy. The statistics of selected transitions for the four site and the seven site models, are explored in the remaining part of this section.

Four interacting sites exhibit the combination of the features of the two site and the three site models, as illustrated in Fig. 2. Figure 2 (b) exhibits the new feature that typical trajectories ($s = 0$) have different statistics at different temperatures: anti-bunching at higher temperatures and bunching at lower temperatures. In Fig. 2 (c) the trajectories remain anti-bunched over the computed temperature range, similar to the results of the two site model.

We consider all possible pairwise jumps of the four site model and conclude that *for sufficiently high activations all trajectories become anti-bunched and approach a temperature independent but transition dependent value of the Mandel parameter.* For sufficiently low activations, on the other hand, all trajectories approach the Poissonian limit. Trajectories that undergo a statistical crossover, approach the Poissonian limit from the positive values of $Q(s)$; those that do not, remain in the $Q(s) < 0$ portion. Approach to the Poissonian limit from the $Q(s) > 0$ side, is therefore a sufficient




Figure 1. a) Excitation states with only two sites present and the corresponding schematics of the interacting sites. The circle containing both sites indicates that they are strongly interacting. b) Excitation states in the three site model. The three site model consists of an strongly interacting pair, weakly interacting with a third site. For each case the statistics of jumps between the excitons associated with sites 1 and 2 are computed. c) $\theta(s)$ for the two site model, d) and the three site model. e) $Q(s)$ for the two site, f) and the three site model. The site energies are $\epsilon_1 = 200 \text{ cm}^{-1}$, $\epsilon_2 = 320 \text{ cm}^{-1}$, $\epsilon_3 = 0 \text{ cm}^{-1}$ and the electronic couplings are $J_{12} = -87.7 \text{ cm}^{-1}$, $J_{13} = 5.5 \text{ cm}^{-1}$, $J_{23} = 30.8 \text{ cm}^{-1}$ [24].

indicator of the existence of statistical crossovers. These conclusions are also observed to be valid for larger systems.

Finally, statistics of selected jumps from the seven site system of the FMO complex are illustrated in Fig. 3. We observe that fast transitions are more likely to show statistical crossovers. The transfer rate between two exciton states separated in energy by ω is given by $\Gamma(\omega) = \gamma(\omega) \sum_m |c_m(\alpha)|^2 |c_m(\alpha')|^2$. In this system, the relative magnitude of the transfer rates tends to be dominated by the interference contribution $\sum_m |c_m(\alpha)|^2 |c_m(\alpha')|^2$, as the variation of $\gamma(\omega)$ over the energy range of interest is small. In other words, *dynamical crossovers are observed in trajectories accounting for jumps between strongly interfering eigenstates*. The dynamics of the rare trajectories are therefore sensitively dependent on the distribution of the electronic couplings and the degree of delocalization and interference. For exciton states that are close in energy,




Figure 2. a) Exciton states for the four site model, and the corresponding schematics of the interacting sites. The four site model consist of two groups of strongly interacting dimers. b) and c) Mandel parameter versus the activation parameter for selected downward jumps in the four site model. The additional site energy is $\epsilon_4 = 110\text{ cm}^{-1}$, and the additional electronic couplings are $J_{14} = -5.9\text{ cm}^{-1}$, $J_{24} = 8.2\text{ cm}^{-1}$, $J_{34} = -53.5\text{ cm}^{-1}$ [24].

$\gamma(\omega)$ is large, resulting in fast transfer. Statistical crossovers are therefore also observed in this regime, despite the small exciton interference. Example of this behaviour can be seen in Fig. 3 (a).

There are important differences between the statistics of quantum jumps in EET, and those of the familiar problems of electron and photon counting that merit further elaboration. Electron counting is fundamentally different from photodetection in that electrons are not destroyed, but are simply counted [4]. Quantum jumps in EET share certain characteristics of each example: they are photon-like as a jump corresponds to transitions between eigenstates; and electron-like as the detection scheme is assumed to be a simple counter that does not interference with the state of the system. Unlike photon counting, transitions between eigenstates are thermally activated (they are not accompanied by emission of photons), and unlike electron counting, the jumps do not generate a current. The insights of photon statistics and current fluctuations should therefore only be applied with caution.

We conclude by noting that the importance of the Markovian assumption in the present treatment poses a difficulty for many systems exhibiting EET whose environmental interactions are strongly non-Markovian. The formalism can, however, be applied to model certain types of non-Markovianity by extending the Hilbert space of the system Hamiltonian to contain the environmental modes that capture the strong non-Markovian features [26, 27].




Figure 3. a) to d) Mandel parameter for selected downward transitions. Colours indicate the temperature of the bath: red (300 K), green (200 K) and blue (100 K). e) Exciton states of the 7 site system and the corresponding schematics of the interacting sites. The seven site system consists of three strongly interacting dimers, coupled weakly to a seventh site.

5. Conclusion

Markovian master equations are an established method of modelling the dynamics of many systems exhibiting transport properties. These equations are primarily used to compute the ensemble averages, but in fact contain more information pertaining to higher order statistical moments that cannot be gleaned unless an unravelling formalism is first implemented. By introducing an artificial coupling field, or an activation parameter, the s -ensemble theory allows the unravelling formalism to be extended to quantum trajectories that have been perturbed out of equilibrium. Such an unravelling formalism has been applied in the present article to analyze the statistics of Markovian excitation transport.

The non-equilibrium steady-state of a system undergoing EET exhibits surprisingly rich statistics: The features that are often associated with quantum correlations such as bunching and anti-bunching can be observed in the steady-state statistics of the thermally-activated quantum jump trajectories. System size also plays a key role in the non-equilibrium statistics as systems of two and three sites reveal qualitatively different correlations: while in the former the jumps are always sub-Poissonian, in the latter they can undergo statistical crossovers. Transitions in larger systems show a combination of these two characteristic behaviours. Moreover, statistical crossovers are found to be activated by the electronic coherences: trajectories that account for jumps between

two strongly interfering eigenstates are most likely to exhibit statistical crossovers. The structure of the spectral density can also influence the statistics by promoting electronic transitions of particular frequencies [28].

We leave the matter of the potential implications of the observed transitions to excitation transport as an intriguing open question. As the quantum jumps are not directly observable, their statistics cannot be directly probed. However, if the statistics can be mapped to an observable of the system, their dependence on the structure of the bath suggests that the observable can be used as a mean to probe the system-bath interactions and non-equilibrium perturbations of the environment.

6. Acknowledgment

J. P. Garrahan is gratefully acknowledged for numerous stimulating discussions. H. H. also acknowledges the hospitality of University College London. The Engineering and Physical Research Council (EPSRC) of the UK and the Natural Sciences and Engineering Research Council of Canada (NSERC) are also thanked for financial support.

References

- [1] L. S. Levitov and G. B. Lesovik. *JETP Lett.*, 58:230, 1993.
- [2] L. S. Levitov, H. W. Lee, and G. B. Lesovik. *J. Maths. Phys.*, 37:10, 1996.
- [3] Ya. M. Blanter and M. Buettiker. Shot noise in mesoscopic conductors. *Phys. Rep.*, 336:1, 2000.
- [4] Y. V. Nazarov. *Quantum Noise in Mesoscopic physics*. Kluwer academic publishers, Dordrecht, The Netherlands, 2002.
- [5] D. A. Bagrets and Yu. V. Nazarov. Full counting statistics of charge transfer in coulomb blockade systems. *Phys. Rev. B*, 67:085316, 2003.
- [6] W. Belzig. Full counting statistics of super-poissonian shot noise in multilevel quantum dots. *Phys. Rev. B*, 71:161301, 2005.
- [7] A. Braggio, J. Konig, and R. Fazio. Full counting statistics in strongly interacting systems: Non-markovian effects. *Phys. Rev. Lett.*, 96:026805, 2006.
- [8] S. Gustavsson, R. Leturcq, B. Simovic, R. Schleser, T. Ihn, P. Studerus, and K. Ensslin. Counting statistics of single electron transport in a quantum dot. *Phys. Rev. Lett.*, 96:076605, 2006.
- [9] Y. J. Zheng and F. H. L. Brown. Single-molecule photon counting statistics via generalized optical bloch equations. *Phys. Rev. Lett.*, 90:238305, 2003.
- [10] Y. J. Zheng and F. H. L. Brown. Photon emission from driven single molecules. *J. Chem. Phys.*, 119:11814, 2003.
- [11] C. Flindt, T. Novotny, and A.-P. Jauho. Full counting statistics of nano-electromechanical systems. *Europhys. Lett.*, 69:475, 2005.
- [12] H.-P. Breuer and F. Pertruccione. *The theory of open quantum systems*. Oxford University Press, New York, 2002.
- [13] J. P. Garrahan and I. Lesanovsky. Thermodynamics of quantum jump trajectories. *Phys. Rev. Lett.*, 104:160601, 2010.
- [14] H. Touchette. The large deviation approach to statistical mechanics. *Physics Reports*, 478:1, 2009.
- [15] A. Budini. Thermodynamics of quantum jump trajectories in systems driven by classical fluctuations. *Phys. Rev. E*, 82:061106, 2010.

- [16] J. P. Garrahan, A. D. Armour, and I. Lesanovsky. Quantum trajectory phase transitions in the micromaser. *Phys. Rev. E*, 84:021115, 2011.
- [17] S. Genway, J. P. Garrahan, I. Lesanovsky, and A. D. Armour. Phase transitions in trajectories of a superconducting single-electron transistor coupled to a resonator. *Phys. Rev. E*, 85:051122, 2012.
- [18] J. Li, Y. Liu, J. Ping, S.-S. Li, X.-Q. Li and Y. Yan . Large-deviation analysis for counting statistics in mesoscopic transport. *Phys. Rev. B*, 84:115319, 2011.
- [19] J. M. Hickey, S. Genway, I. Lesanovsky, and J. P. Garrahan. Thermodynamics of Quadrature Trajectories in Open Quantum Systems. arXiv:1206.5719.
- [20] M. Esposito, U. Harbola, and S. Mukamel. Nonequilibrium fluctuations, fluctuation theorems, and counting statistics in quantum systems. *Rev. Mod. Phys.*, 81:1665, 2009.
- [21] A. Budini. Large deviations of ergodic counting processes: A statistical mechanics approach. *Phys. Rev. E*, 84:011141, 2011.
- [22] M. B. Plenio and P. L. Knight. The quantum jump approach to dissipative dynamics in quantum optics. *Rev. Mol. Phys.*, 70:101–144, 1998.
- [23] M. Mohseni, P. Rebentrost, S. Lloyd, and A. Aspuru-Guzik. Environment-assisted quantum walks in photosynthetic energy transfer. *J. Chem. Phys.*, 129:174106, 2008.
- [24] J. Adolphs and T. Renger. How proteins trigger excitation energy transfer in the FMO complex of green sulfur bacteria. *Biophys. J.*, 91:2778–2797, 2006.
- [25] F. Fassioli, A. Nazir, and A. Olaya-Castro. Distribution of entanglement in light-harvesting complexes and their quantum efficiency. *New J. Phys.*, 12:8, 2010.
- [26] L. Mazzola, S. Maniscalco, J. Piilo, K.-A. Suominen, and B. M. Garraway. Pseudomodes as an effective description of memory: Non-markovian dynamics of two-state systems in structured reservoirs. *Phys. Rev. A*, 80:012104, 2009.
- [27] H.-P. Breuer. Genuine quantum trajectories for non-markovian processes. *Phys. Rev. A*, 70:012106, 2004.
- [28] H. Hossein-Nejad, C. Curutchet, A. Kubica, and G. D. Scholes. Delocalization-enhanced long-range energy transfer between cryptophyte algae PE545 antenna proteins. *J. Phys. Chem. B.*, 115:5243–53, 2011.